

- B5
amub
- (b) cultivating the cells under a selection pressure;
 - (c) selecting or screening for one or more transformants expressing a desired characteristic; and
 - (d) isolating the transformant(s) of interest.
-

B6
Sub 3

(Amended) The method of claim 1, wherein the polynucleotide sequence further comprises a control sequences.

4. (Amended.) The method according to claim 1, wherein the polynucleotide sequence of interest encodes a hormone, an enzyme, a receptor or a portion thereof, an antibody or a portion thereof, or a reporter, or a regulatory protein.
-

- B7
7. (Amended) The method according to claim 3, wherein the control sequence is an enhancer sequence, a leader sequence, a polyadenylation sequence, a propeptide sequence, a promoter, a replication initiation sequence, a signal sequence, a transcriptional terminator or a translational terminator.
-

- B8
9. (Amended) The method according to claim 1, wherein the selection marker polynucleotide sequence is selected from the group of genes which encode a product which is responsible for one of the following: resistance to biocide or viral toxicity, resistance to heavy metal toxicity, prototrophy to auxotrophs.
-

- B9
11. (Amended.) The method of claim 9, wherein the selection marker polynucleotide sequence is a gene selected from the group consisting of *argB* (ornithine carbamoyltransferase), *amdS* (acetamidase), *bar* (phos-hinotricin acetyltransferase), *hemA* (5-aminolevulinate synthase), *hemB* (porphobilinogen synthase), *hygB* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *prn* (proline permease), *pyrG* (orotidine-5'-phosphate decarboxylase), *pyroA*, *riboB*, *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase).
-

12. (Amended) The method of claim 1, wherein the replication initiating sequence is a nucleic
- 3 -

acid sequence selected from the group consisting of:

- B9
amended
- (a) a replication initiating sequence having at least 80% identity with the nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:2, as determined using the GAP computer program with a GAP creation penalty of 5.0 and GAP extension penalty of 0.3, and is capable of initiating replication; and
 - (b) a replication initiating sequence which hybridises under low stringency conditions with (i) the nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:2, or (ii) the respective complementary strands, wherein the low stringency conditions are defined by prehybridization and hybridization at 42°C in 5x SSPE, 0.3% SDS, 200 mg/ml sheared and denatured salmon sperm DNA, and 25% formamide, and wash conditions are defined at 50°C for 30 minutes in 2x SSC, 0.2% SDS.

Sub
C2

13. (Amended) The method of claim 12, wherein the nucleic acid sequence has at least 80% identity with the nucleic acid sequence of SEQ ID NO:1 or SEQ ID NO:2, as determined using the GAP computer program with a GAP creation penalty of 5.0 and GAP extension penalty of 0.3.

14. (Amended.) The method of claim 12, wherein the replication initiating polynucleotide sequence is obtained from a filamentous fungal cell.

B10
Sub
C3

17. (Twice amended.) The method of claim 12, wherein the replication initiating polynucleotide sequence has the nucleic acid sequence set forth in SEQ ID NO:1 or SEQ ID NO:2.

18. (Amended.) The method of claim 2, wherein the polynucleotide sequence of interest was created by mutagenesis, by random mutagenesis, by use of a physical or chemical mutagenizing agent, by use of a doped oligonucleotide, by DNA shuffling, by subjecting the nucleic acid sequence to PCR generated mutagenesis, or by use of any combination thereof.

Please add new claim 30, as follows:

B11
Sub
C4

30. (New.) The method of claim 1, wherein the polynucleotide sequence of interest is a control

B¹¹
Contd

sequences.
